

AD-A166 917


GAMBLE-II IMPLoding SODIUM PLASMA I CALIBRATION OF THE
HELIUM-LIKE RESONANCE (U) NAVAL RESEARCH LAB WASHINGTON
DC J DAVIS ET AL 10 APR 86 NRL-MR-5765


1/1

UNCLASSIFIED

F/G 20/5

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

2

NRL Memorandum Report 5765

Gamble-II Imploding Sodium Plasma

I. Calibration of the Heliumlike Resonance Line as a Pump Source and Detection of Fluorescence in Neon

J. DAVIS, J. E. ROGERSON AND J. P. APRUZESE

*Plasma Radiation Branch
Plasma Physics Division*

April 10, 1986

This research was sponsored by the Defense Nuclear Agency under Subtask QIEQMXLA, work unit 00006 and work unit title "XRL Source."



NAVAL RESEARCH LABORATORY
Washington, D.C.

Approved for public release; distribution unlimited.

DTIC
ELECTE
MAY 06 1986
S E D

86 5 5 068

AD-A166 917

DTIC FILE COPY

SECURITY CLASSIFICATION OF THIS PAGE

REPORT DOCUMENTATION PAGE

1a. REPORT SECURITY CLASSIFICATION UNCLASSIFIED			1b. RESTRICTIVE MARKINGS		
2a. SECURITY CLASSIFICATION AUTHORITY			3. DISTRIBUTION / AVAILABILITY OF REPORT		
2b. DECLASSIFICATION / DOWNGRADING SCHEDULE			Approved for public release; distribution unlimited.		
4. PERFORMING ORGANIZATION REPORT NUMBER(S) NRL Memorandum Report 5765			5. MONITORING ORGANIZATION REPORT NUMBER(S)		
6a. NAME OF PERFORMING ORGANIZATION Naval Research Laboratory		6b. OFFICE SYMBOL (if applicable) Code 4720		7a. NAME OF MONITORING ORGANIZATION	
6c. ADDRESS (City, State, and ZIP Code) Washington, DC 20375-5000			7b. ADDRESS (City, State, and ZIP Code)		
8a. NAME OF FUNDING / SPONSORING ORGANIZATION Defense Nuclear Agency		8b. OFFICE SYMBOL (if applicable)		9. PROCUREMENT INSTRUMENT IDENTIFICATION NUMBER	
8c. ADDRESS (City, State, and ZIP Code) Alexandria, VA 22310			10. SOURCE OF FUNDING NUMBERS		
			PROGRAM ELEMENT NO. 62715H	PROJECT NO.	TASK NO.
			WORK UNIT ACCESSION NO. DN155-166		
11. TITLE (Include Security Classification) Gamble-II Imploding Sodium Plasma — I. Calibration of the Heliumlike Resonance Line as a Pump Source and Detection of Fluorescence in Neon					
12. PERSONAL AUTHOR(S) Davis, J., Rogerson, J. E. and Apruzese, J. P.					
13a. TYPE OF REPORT Interim		13b. TIME COVERED FROM 10/85 TO 1/86		14. DATE OF REPORT (Year, Month, Day) 1986 April 10	
15. PAGE COUNT 33					
16. SUPPLEMENTARY NOTATION This research was sponsored by the Defense Nuclear Agency under Subtask QIEQMXLA, work unit 00006 and work unit title "XRL Source."					
17. COSATI CODES			18. SUBJECT TERMS (Continue on reverse if necessary and identify by block number)		
FIELD	GROUP	SUB-GROUP	→ Imploding Resonance line X-ray laser ← Sodium plasma Pump		
19. ABSTRACT (Continue on reverse if necessary and identify by block number) Numerical simulations are performed describing the behavior and properties of an imploding cylindrical annular sodium plasma. Special attention is devoted toward calibrating the power radiated in the heliumlike resonance line for application as a pump source in x-ray laser experiments. Detection of fluorescence in a nearby neon plasma is also discussed. <i>keywords</i>					
20. DISTRIBUTION / AVAILABILITY OF ABSTRACT <input checked="" type="checkbox"/> UNCLASSIFIED/UNLIMITED <input type="checkbox"/> SAME AS RPT. <input type="checkbox"/> DTIC USERS			21. ABSTRACT SECURITY CLASSIFICATION UNCLASSIFIED		
22a. NAME OF RESPONSIBLE INDIVIDUAL Jack Davis			22b. TELEPHONE (Include Area Code) (202) 767-3278		22c. OFFICE SYMBOL Code 4720

DD FORM 1473, 84 MAR

63 APR edition may be used until exhausted.
All other editions are obsolete.

SECURITY CLASSIFICATION OF THIS PAGE

CONTENTS

I.	INTRODUCTION	1
II.	RESULTS AND DISCUSSION	2
	ACKNOWLEDGMENTS	7
	REFERENCES	19

Accession For	
NTIS GRA&I	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By _____	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



GAMBLE-II IMPLoding SODIUM PLASMA
I. CALIBRATION OF THE HELIUMLIKE RESONANCE LINE AS A
PUMP SOURCE AND DETECTION OF FLUORESCENCE IN NEON

I. Introduction

A variety of schemes have been proposed suggesting ways of creating a population inversion and gain coefficients in excess of unity in the x-ray region. One attractive scheme involves the flashlamp concept where the radiated flux from one plasma pumps another plasma to create conditions favorable to the emission of coherent radiation. Of the many photopumped schemes based on line coincidence, the Na-Ne scheme is particularly interesting because the lines match to two parts in 10^4 at a wavelength of 11Å. In particular, the NaX $1s^2\ ^1S_0-1s2p\ ^1P_1$ line at 11.0027Å pumps the NeIX $1s^2\ ^1S_0-1s4p\ ^1P_1$ line at 11.0003Å. The strongest lasing lines in the pumped neon system are the $2p\ ^1P_1-3d\ ^1D_2$, $2p\ ^1P_1-4d\ ^1D_2$, and $3d\ ^1D_2-4f\ ^1F_3$. This scheme is particularly interesting because: (1) both ions are heliumlike - a closed shell configuration exists over a large parameter space of temperature and density, (2) the line coincidence is almost perfect - two parts in 10^4 - which is about a Doppler width, and (3) both ions can be produced easily in the laboratory by Z-pinch plasmas driven by pulse power generators and laser plasmas. A comprehensive discussion on the kinetics and the plasma environment required for lasing to occur has been presented in two earlier papers.^{1,2} The details of the calculations can be found there. In this paper we will focus on the sodium pump source and determine the feasibility of producing a heliumlike sodium plasma radiating an intense flux in the heliumlike resonance line by means of a current driven gas puff implosion.

The treatment presented here is based on the implosion dynamics of a cylindrical annular gas puff plasma consisting initially of sodium vapor. The implosion is driven by a high current discharge typical of the GAMBLE

II pulse power generator. The theory describing the implosion is based on a dynamic pinch model self-consistently coupled to a non-LTE ionization dynamic and radiation transport model.³ The radiation transport employs a probability of escape method. This provides a self-consistent picture of the temperature, density, size, and level population during the plasmas' evolution. Such an simplified approach ignores any pathological behavior such as instabilities sometimes exhibited by these imploding plasmas. However, the approach adopted here does provide a good starting point for determining the feasibility of producing the radiation flux in the heliumlike resonance line required to pump the $1s^2-1s4p$ level of heliumlike neon leading to a population inversion and gain coefficients in excess of 1cm^{-1} .

II. Results and Discussion

The non-LTE dynamic pinch model has been discussed in considerable detail elsewhere. The reader is referred to Ref. 3 for further particulars. The collisional-radiative model for sodium consists of ground states for all the ionization stages and contains structure up to principal quantum number $n=5$ in the lithium-, helium-, and hydrogen-like ionization stages. The $n=4$ and 5 levels have been averaged and are represented as single levels.

For illustrative purposes, the results of numerical simulations are presented for $30\mu\text{gm/cm}$ of sodium uniformly distributed between the initial inner and outer radii of 0.95 and 1.55 cm, respectively and 4 cm in length. The driving current waveform typical of the GAMBLE II generator is shown in Fig. 1. The current reaches a peak value of 1.25 megamp in about 70 nsec and then decays to about 0.45 megamp in about 50 nsec. The remainder of the current waveform, i.e., beyond about 120 nsec, is for computational expediency. The temporal behavior of the inner (A) and outer (B) radius is shown in Fig. 2. The inner radius collapses and stagnates on axis while the outer radius continues inward until the fluid pressure retards the forward runin motion and the plasma bounces. The final pinch radius is about 1/10 the initial outer radius. The variation of velocity with time is shown in Fig. 3, attaining a peak value of $23\text{ cm}/\mu\text{sec}$ at about

120 nsec which is well after peak current. This behavior is characteristic of the GAMBLE II current driven implosions. The temperature and density time histories are shown in Figs. 4 and 5, respectively. The temperature profile peaks at about 380 eV while the density profile peaks at 2×10^{19} ions/cm³, both profiles maximizing around 140 nsec. These values are similar to those obtained for neon⁴ which is not too surprising since sodium is just one atomic number away from neon. The total radiative yield as a function of time is shown in Fig. 6 and is about 4.7 kilojoules. The power radiated (in Watts) is presented as a function of time (nsec) in Fig. 7 for radiation in energy bins labelled K- and L-lines. The K-lines refer to radiation originating from transitions equal to or above 1 KeV and L-lines representing those transitions below 1 KeV. Within the context of this study the L-lines refer strictly to transitions with energies less than 1 keV emanating from the Li-, He-, and H-like ionization stages only. The L-lines appear first and linger longer than the K-lines but the K-lines reach a peak value greater than the corresponding peak value attained in the L-lines. Also, in the case of argon, the K-lines peak occurs just before the peak in the L-lines.⁵ The reasons for this have been discussed in detail elsewhere and need not concern us here.³ The emission spectrum at 141 nsec is given as a function of energy (KeV) in Fig. 8. The line intensities are fairly well distributed with peaks occurring in the K-shell above 1KeV and in a band of lines around 250 eV. This result, is to some extent, model dependent in the sense that an extended sodium model containing more transitions would provide a richer and possibly modified spectrum than the one presented in Fig. 8. Certainly, the total power radiated would increase with an extended model. Since our primary concern was to theoretically calibrate the flux emitted from the heliumlike resonance line, i.e., the flux in the $1s^2 \ ^1S - 1s2p \ ^1P$ transition, we present in Fig. 9 the power radiated in this line as a function of time. At or near the pinch the power radiated is about 8×10^{10} watts. This is in very good agreement with the peak value of 5×10^{10} watts obtained experimentally with neon on the GAMBLE II facility.⁶

In order to determine the behavior and magnitude of the radiation flux from the He _{α} -like resonance line for a variety of conditions, numerical simulations were carried out for B = 1.55 cm and A = 0.95 cm (outer and inner radii, respectively) as a function of M/l. These results are

summarized in Fig. 10. Fairly impressive yields can be obtained for the above conditions with a peak value of 0.08 terrawatts when M/l is roughly between 25 and 30 $\mu\text{gm/cm}$. For higher and lower values of M/l the yield drops precipitously. For higher values of M/l , i.e., higher than, say, 40 $\mu\text{gm/cm}$, the load becomes too massive to achieve large values of runin velocity and subsequently is inefficient at converting kinetic energy to thermal energy at high temperatures commensurate with the heliumlike ionization stage. On the other hand, for loads of small mass, the conversion of runin kinetic energy to thermal energy causes overheating and burn-through of the heliumlike ionization stage leading to a reduction in the emission from the heliumlike ions, provided, of course, the plasma is thermal.

Similarly, for fixed values of $M/l=30\mu\text{gm/cm}$ and $B=1.55\text{ cm}$, results were obtained for the power radiated in the heliumlike resonance line by varying the inner radius. These results are displayed in Fig. 11 as a function of ΔR , the difference between the outer and inner radius. The total power radiated in the He_α -like resonance line over the range of values considered is relatively insensitive to ΔR , having a value of 0.08 terrawatts at $\Delta R=0.15\text{ cm}$ and falling to 0.07 terrawatts at $\Delta R=0.50\text{ cm}$, primarily because for the value of $M/l=30\mu\text{gm/cm}$, the load is efficiently coupled to the generator in terms of generating K-shell radiation.

From these calculations it is apparent that if it is experimentally feasible to implode a cylindrical annular sodium plasma under the right set of conditions, then it is theoretically possible to attain a reasonably potent pump source from the heliumlike resonance line of the order of 0.08 terrawatts at 11.0027Å. The question confronting us now is whether this pump source is adequate to pump the heliumlike neon $1s^2\ ^1S-1s4p\ ^1P$ transition and create a population inversion in the upper state that may subsequently lead to gain coefficients in excess of unity. To answer this question we assume a neon plasma to be 1 cm away from the sodium pump source. The area of a cylinder centered on the pump 1 cm away from the pump is 25.1 cm^2 . Also, the line width of the $1s^2\ ^1S-1s2p\ ^1P$ line is about $3 \times 10^{14}\text{ Hz}$, hence, the pump flux is

$$\text{Pump Flux} = \frac{8 \times 10^{10} \text{ watts} \cdot 10^7 \text{ erg/joule}}{(3 \times 10^{14} \text{ Hz}) (25.1 \text{ cm}^2)}$$

$$= 10^2 \frac{\text{ergs}}{\text{cm}^2 \text{-sec-Hz}}$$

This is roughly the equivalent of a 125 eV blackbody source. From our earlier calculations this would lead to a gain of unity in a cool neon plasma just ionized to the heliumlike stage at 1 cm from the sodium plasma. Even if the neon cannot be placed at a distance of 1 cm from the sodium, fluorescence of the neon would easily be detectable and would represent a major step forward.

To further explore detection of fluorescence in heliumlike neon pumped by heliumlike sodium in Gamble II, assume that a cool (65-100 eV), low-density ($N_I < 10^{18} \text{ cm}^{-3}$) neon plasma, barely ionized to the heliumlike stage, can be placed within 2 cm or so of the sodium pinch pump in Gamble-II. The kinetics of the levels of such a heliumlike neon plasma would be dominated by the sodium pump radiation to the $1s4p^1P$ level and resulting cascade of radiative decays. In this case, a simple model can yield reasonable estimates of what the fluorescence-induced heliumlike neon resonance line ratios would be.

LEVEL MODEL

- 1 = $1s^2 \ ^1S_0$ ground state in heliumlike neon
- 2 = $1s2p^1P$
- 3 = $1s3l$ singlets
- 4 = $1s4l$ singlets

Definitions

P = pump rate of heliumlike neon $1s4l$ singlet level by sodium (sec^{-1}), per ground state ion

$A_{43}, A_{42}, A_{41}, A_{32}, A_{31}, A_{21}$, = spontaneous decay rates for the indicated transitions (Einstein A's), in sec^{-1}

f_1 = ground state fraction

(f_2, f_3, f_4) = excited state fractions

Equations for Steady State

<u>CREATION RATE</u>	=	<u>DESTRUCTION RATE</u>
Level 4: $f_1 P$	=	$f_4(A_{43} + A_{42} + A_{41})$
Level 3: $f_4 A_{43}$	=	$f_3(A_{32} + A_{31})$
Level 2: $f_4 A_{42} + f_3 A_{32}$	=	$f_2(A_{21})$

Decay Rates

$$A_{42} = 4.6 \times 10^{10} \text{ sec}^{-1}, A_{43} = 6.0 \times 10^{10} \text{ sec}^{-1}, A_{41} = 1.9 \times 10^{11} \text{ sec}^{-1}$$
$$A_{32} = 2.6 \times 10^{11} \text{ sec}^{-1}, A_{31} = 8.3 \times 10^{11} \text{ sec}^{-1}, A_{21} = 8.9 \times 10^{12} \text{ sec}^{-1}$$

Simple Algebra Yields:

$$f_4/f_1 = 3.38 \times 10^{-12} P$$

$$f_3/f_1 = 1.86 \times 10^{-13} P$$

$$f_2/f_1 = 2.29 \times 10^{-14} P$$

Therefore

$$f_4, f_3, f_2 \text{ are in the ratio } 1.0 : 0.055 : 0.00678.$$

The line intensities (optically thin) are in the ratio

$$f_4 A_{41} : f_3 A_{31} : f_2 A_{21} = 1.0 : 0.24 : 0.32.$$

Therefore, the 1→4 resonance line will be three times as intense as the principal resonance line at 13.45 Å. In case the 1→4 radiation cannot be experimentally separated from the pumping sodium radiation, the 1→3/1→2 ratio of 0.75 is also far above what would be expected (0.1-0.2) for an isolated neon plasma, and would indicate fluorescence and pumping.

ACKNOWLEDGMENTS

This work was supported in part by SDIO, through DNA, and ONR. We would also like to express our thanks to Dr. D. Duston for developing the rate tables and Dr. P. Kepple for making them available. Also, we thank Chris Agritellis for his assistance in making the SIMPLODE code operational on the VAX computer and to Dr. F. C. Young for his comments on the manuscript.




Fig. 1. Current waveform as a function of time.




Fig. 2. Radii as a function of time.




Fig. 3. Velocity as a function of time.




Fig. 4. Temperature as a function of time.




Fig. 5. Ion density, reduced by 10^{10} , as a function of time.




Fig. 6. Total radiative yield as a function of time.




Fig. 7. Power radiated in L-and K-lines as a function of time.




Fig. 8. Radiated emission spectrum as a function of energy.




Fig. 9. Power radiated in heliumlike sodium resonance line as a function of time.




Fig. 10. Power radiated in heliumlike sodium resonance line as a function of M/l for $\Delta R = 0.60$ cm.




Fig. 11. Power radiated in heliumlike sodium resonance line as a function of ΔR for $M/\lambda = 30 \text{ } \mu\text{gm/cm}$.

References

1. F. L. Cochran, J. Davis, and J. P. Apruzese, J. Appl. Phys. 57, 27 (1985).
2. J. P. Apruzese and J. Davis, Phys. Rev. A 31, 2976 (1985).
3. J. Davis, C. Agritellis, and D. Duston, NRL Memo Report 5615 (1985).
Simplode: An Imploding Gas Puff Plasma Model I. Neon (ADA157456)
4. S. J. Stephanakis, J. P. Apruzese, P. G. Burkhalter, J. Davis, R. A. Meger, S. W. McDonald, C. G. Mehlman, P. F. Ottinger and F. C. Young, J. Appl. Phys. Lett. (to be published).
5. F. C. Young, S. J. Stephanakis, V. E. Scherrer, J. R. Boller, P. F. Ottinger and S. W. McDonald, Bull. Am. Phys. Soc. 30, 1389 (1985).
6. S. J. Stephanakis, S. W. McDonald, P. F. Ottinger, J. R. Boller, V. E. Scherrer and F. C. Young, Bull. Am. Phys. Soc. 30, 1389 (1985).

DISTRIBUTION LIST

Assistant to the Secretary of Defense Atomic Energy Washington, D.C. 20301 ATTN: Executive Assistant	1 Copy
Defense Technical Information Center Cameron Station 5010 Duke Street Alexandria, Va 22314	2 copies
Director Defense Intelligence Agency Washington, D.C. 20301 ATTN: DT-1B R. Rubenstein	1 Copy
Director Defense Nuclear Agency Washington, D.C. 20305 ATTN: DDST ATTN: TITL ATTN: RAEV ATTN: STVI	1 copy 4 copies 1 copy 1 copy
Commander Field Command Defense Nuclear Agency Kirtland AFB, New Mexico 87115 ATTN: FCPR	1 Copy
Chief Field Command Livermore Division Department of Defense P.O. Box 808 Livermore, CA 94550 ATTN: FCPRL	1 Copy
Director Joint Strat TGT Planning Staff Offutt AFB Omaha, Nebraska 68113 ATTN: JLKS	1 Copy
Undersecretary of Defense for RSCH and ENGRG Department of Defense Washington, D.C. 20301 ATTN: Strategic and Space Systems (OS)	1 Copy

Deputy Chief of Staff for RSCH DEV and ACQ
Department of the Army
Washington, D.C. 20301
ATTN: DAMA-CSS-N

1 Copy

Commander
Harry Diamond Laboratories
Department of the Army
2800 Powder Mill Road
Adelphi, MD 20783
ATTN: DELHD-N-NP
ATTN: DELHD-R J. Rosado
ATTN: DELHD-TA-L (Tech. Lib.)

1 copy each

U.S. Army Missile Command
Redstone Scientific Information Center
Attn: DRSMI-RPRD (Documents)
Redstone Arsenal, Alabama 35809

3 Copies

Commander
U.S. Army Missile Command
Redstone Arsenal, Alabama 35898
ATTN: DRCPM-PE-EA

1 copy

Commander
U.S. Army Nuclear and Chemical Agency
7500 Backlick Road
Building 2073
Springfield, VA 22150
ATTN: Library

1 copy

Commander
Naval Intelligence Support Center
4301 Suitland Road, Bldg. 5
Washington, D.C. 20390
ATTN: NISC-45

1 Copy

Commander
Naval Weapons Center
China Lake, California 93555
ATTN: Code 233 (Tech. Lib.)

1 Copy

Officer in Charge
White Oak Laboratory
Naval Surface Weapons Center
Silver Spring, Md. 20910
ATTN: Code R40
ATTN: Code F31

1 Copy each

Air Force Weapons Laboratory Kirtland AFB, New Mexico 87117 ATTN: SUL ATTN: CA ATTN: APL ATTN: Lt. Col Generosa	1 Copy each
Deputy Chief of Staff Research, Development and Accounting Department of the Air Force Washington, D. C. 20330 ATTN: AFRDQSM	1 Copy
Commander U.S. Army Test and Evaluation Command Aberdeen Proving Ground, MD 21005 ATTN: DRSTE-EL	1 Copy
AVCO Research and Systems Group 201 Lowell Street Wilmington, MA 01887 ATTN: Library A830	1 Copy
BDM Corporation 7915 Jones Branch Drive McLean, Virginia 22101 ATTN: Corporate Library	1 Copy
Berkeley Research Associates P.O. Box 983 Berkeley, CA 94701 ATTN: Dr. Joseph Workman	1 Copy
Berkeley Research Associates P.O. Box 852 5532 Hempstead Way Springfield, VA 22151 ATTN: Dr. Joseph Orens	1 Copy each
Boeing Company P. O. Box 3707 Seattle, WA 98134 ATTN: Aerospace Library	1 Copy
The Dikewood Corporation 1613 University Bldv., N.E. Albuquerque, New Mexico 87110 ATTN: L. Wayne Davis	1 Copy

General Electric Company
Space Division
Valley Forge Space Center
P. O. Box 8555
Philadelphia, PA 19101
ATTN: J. Peden

1 Copy

General Electric Company - Tempo
Center for Advanced Studies
816 State Street
P.O. Drawer QQ
Santa Barbara, CA 93102
ATTN: DASIAC

1 Copy

Institute for Defense Analyses
1801 N. Beauregard St.
Alexandria, VA 22311
ATTN: Classified Library

1 Copy

IRT Corporation
P.O. Box 81087
San Diego, CA 92138
ATTN: R. Mertz

1 Copy

JAYCOR
11011 Forreyane Rd.
P.O. Box 85154
San Diego, CA 92138
ATTN: E. Wenaas
F. Felbar

1 Copy

JAYCOR
205 S. Whiting Street, Suite 500
Alexandria, VA 22304
ATTN: R. Sullivan

1 Copy

KAMAN Sciences Corp.
P. O. Box 7463
Colorado Springs, CO 80933
ATTN: Library

1 copy each

Lawrence Livermore National Laboratory
University of California
P.O. Box 808
Livermore, California 94550
Attn: DOC CDN for L-153
Attn: DOC CDN for L-47 L. Wouters
Attn: DOC CDN for Tech. Infor. Dept. Lib.

1 copy each

Lockheed Missiles and Space Co., Inc.
P. O. Box 504
Sunnyvale, CA 94086
Attn: S. Taimlty
Attn: J.D. Weisner

1, copy each

Lockheed Missiles and Space Co., Inc. 3251 Hanover Street Palo Alto, CA 94304 Attn: J. Perez	1 Copy
Maxwell Laboratory, Inc. 9244 Balboa Avenue San Diego, CA 92123 ATTN: A. Kolb ATTN: M. Montgomery ATTN: J. Shannon	1 Copy each
McDonnell Douglas Corp. 5301 Bolsa Avenue Huntington Beach, CA 92647 ATTN: S. Schneider	1 Copy
Mission Research Corp. P. O. Drawer 719 Santa Barbara, CA 93102 ATTN: C. Longmire ATTN: W. Hart	1 Copy each
Mission Research Corp.-San Diego 5434 Ruffin Rd. San Diego, California 92123 ATTN: Victor J. Van Lint	1 Copy
Northrop Corporation Northrop Research and Technology Center 1 Research Park Palos Verdes Peninsula, CA 90274 ATTN: Library	1 Copy
Northrop Corporation Electronic Division 2301 120th Street Hawthorne, CA 90250 ATTN: V. Damarting	1 Copy
Physics International Company 2700 Merced Street San Leandro, CA 94577 ATTN: M. Krishnan ATTN: C. Gilman ATTN: S. Wong	1 Copy each
R and D Associates P.O. Box 9695 Marina Del Rey, CA 90291 ATTN: W. Graham, Jr. ATTN: P. Haas	1 Copy each

Sandia National Laboratories P.O. Box 5800 Albuquerque, New Mexico 87115 ATTN: Doc Con For 3141 ATTN: D. McDaniel ATTN: P. VanDevender ATTN: K. Matzen, Code 4247	1 copy each
Science Applications, Inc. P. O. Box 2351 La Jolla, CA 92038 ATTN: R. Beyster	1 copy
Spire Corporation P. O. Box D Bedford, MA 01730 ATTN: R. Little	1 copy
SRI International 333 Ravenswood Avenue Menlo Park, CA 94025 ATTN: S. Dairiki	1 copy
S-CUBED P. O. Box 1620 La Jolla, CA 92038 ATTN: A. Wilson	1 copy
Director Strategic Defense Initiative Organization 1717 H Street Pentagon 20301-7100 ATTN: DE Lt. Col Richard Gullickson/DEO IST Dr. Dwight Duston ATTN: IST Dr. J. Ionson	1 copy each
Texas Tech University P.O. Box 5404 North College Station Lubbock, TX 79417 ATTN: T. Simpson	1 copy
TRW Defense and Space Systems Group One Space Park Redondo Beach, CA 90278 ATTN: Technical Information Center	1 Copy
University of Buffalo Dept. of Electrical Engineering Attn: Prof. J. Sargent 3435 Main Street Buffalo, New York 14214	1 Copy

Manfred L. Bitter
Plasma Physics Laboratory
Princeton University
Princeton, NJ 08540

E. J. T. Burns
Org. 4242
Sandia Laboratories
Albuquerque, NM 87185

George Charatis
KMS Fusion
3621 S. State Road
Ann Arbor, MI 48106

J. Cooper
JILA
University of Colorado
Boulder, Colorado 80302

R. S. Craxton
Lab. for Laser Energetics
U. of Rochester
250 E. River Road
Rochester, NY 14623

Alan Desilva
University of Maryland
Physics Dept.
College Park, MD 20742

Raymond Dukart
Sandia Laboratories
Albuquerque, NM 87185

Reuben Epstein
Lab. for Laser Energetics
University of Rochester
Rochester, NY 14623

Amnon Fisher
Department of Physics
University of California
Irvine, CA 92717

Richard J. Fortner
LLNL
P. O. Box 808 L-401
Livermore, CA 94550

R. Godwin
LANSL MS 420
Los Alamos, NM 87545

Alan Hauer
M. S. 554
Los Alamos National Laboratory
P. O. Box 1663
Los Alamos, NM 87545

Alan L. Hoffman
Mathematical Sciences Northwest
P. O. Box 1887
Bellevue, Washington, 98009

Walter F. Huebner
M. S. 212
Los Alamos National Laboratory
P. O. Box 1663
Los Alamos, NM 87545

Thomas W. Hussey
Org. 4242
Sandia Laboratories
Albuquerque, NM 87185

B. R. Junker
ONR Code 323
800 North Quincy Street
Arlington, VA 22217

R. L. Kauffman
LLNL
P.O. Box 808
Livermore, CA 94550

Keith Matzen
Org. 4247
Sandia Laboratories
Albuquerque, NM 87185

Robert McCrory
University of Rochester
Lab. for Laser Energetics
250 E. River Road
Rochester, NY 14623

David D. Burgess
The Blackett Laboratory
Imperial College
London SW7 2BZ
ENGLAND

C. Deutsch
Laboratoire de Physique des Plasmas
Universite Paris XI
91405 Orsay
FRANCE

H. W. Drawin
Assoc. EURATON-CEA
Sur La Fusion
92260. Fontenay-Aux-Roses
FRANCE

A. H. Gabriel
Space and Astrophysics Division
Appleton Laboratory
Chilton, Didcot, Oxon.
OX11 0QX, UNITED KINGDOM

Frank E. Irons
Western Mining Corp.
55 MacDonald Street
Kalgoorlie, 6430
WESTERN AUSTRALIA

P. Jaegle
Laboratoire de Spectroscopie
Atomique et Ionique
University Paris-Sud
Orsay, FRANCE

Joseph D. Kilkenny
The Blackett Laboratory
Imperial College
London SW7 2BZ
ENGLAND

Hans-Joach I. Kunze
Inst. Experimentalphysik V
Ruhr Universitat
Postfach 2148
4630 Bochum
WEST GERMANY

James G. Lunney
Department of Physics
Queen's University of Belfast
Belfast, Northern Ireland
UNITED KINGDOM

Carl Moser
Centre European de Calcul Atomique
et Moleculaire
Batiment 506
Universite de Paris XI
91405 Orsay Cedex
FRANCE

N. J. Peacock
UKAEA
Culham University
Abingdon, Oxon
OX 143DB, UNITED KINGDOM

G. J. Pert
Dept. of Applied Physics
University of Hull
Hull HU6; 7RX
ENGLAND, UNITED KINGDOM

Steven Rose
A.W.R.E.-E3
Aldermaston, Reading RG7 4PR
UNITED KINGDOM

David Salzmann
Soreq Nuclear Research Center
Yavne 70600
ISRAEL

Dillon H. McDaniel
Div. 4234
Sandia National Laboratories
Albuquerque, NM 87185

Eugene J. McGuire
Org. 4211
Sandia Laboratories
Albuquerque, NM 87185

Mike Montgomery
Maxwell Laboratories
8835 Balboa Avenue
San Diego, CA 92123

H. Pilloff
ONR Code 323
800 N. Quincy St.
Arlington, VA 22217

Dr. Richard H. Pratt
Department of Physics and Astronomy
University of Pittsburgh
Pittsburgh, PA 15260

M. Richardson
U. of Rochester
Lab for Laser Energetics
250 E. River Road
Rochester, NY 14623

Richard L. Schneider
Physics International Inc.
2700 Merced Street
San Leandro, CA 94577

Stanley Skupsky
Laboratory for Laser Energetics
University of Rochester
250 E. River Road
Rochester, NY 14623

T. F. Stratton
LANSL
P. O. Box 1663
Los Alamos, NM 87544

Michael Stroschio
Department of Energy
Washington, D.C. 20545

David M. Woodall
Dept. of Nuclear Engineering
University of New Mexico
Albuquerque, NM 87131

Barukh Yaakobi
Laboratory for Laser Energetics
University of Rochester
250 E. River Road
Rochester, NY 14623

Naval Research Laboratory

Plasma Radiation Branch

Washington, D.C. 20375-5000

Code 4720 - 50 Copies

Code 4700 - 26 Copies

Code 2628 - 20 Copies

Code 1220 - 1 Copy

Director of Research

U.S. Naval Academy

Annapolis, MD 21402 - 1 Copy

END

DTic

5-86